

University of Groningen

Tinnitus

Bartels, Hilke

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2008

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Bartels, H. (2008). *Tinnitus: new insights into pathophysiology, diagnosis and treatment*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).


The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

CHAPTER 5



**The impact of Type D personality on
health-related quality of life and disease
severity in chronic tinnitus patients is
largely mediated by psychological distress**

**H. Bartels
S.S. Pedersen
M.J. Staal
B.F.A.M. van der Laan
J.L. Middel**

Submitted to Psychosomatic research, July 2008

Abstract

Objective

To evaluate 1) the predictive power of psychological distress (i.e., vital exhaustion, anxiety and depression) on HRQoL and tinnitus severity, and 2) whether the indicators of psychological distress mediate the impact of Type D personality on HRQoL and disease severity in chronic tinnitus patients.

Material and methods

265 consecutive tinnitus patients completed self-report questionnaires. Anxiety and depression were assessed with the the Hospital Anxiety and Depression Scale, vital exhaustion with the Maastricht Questionnaire, Type D with the DS14, and quality of life with the Short Form Health Survey 36 and tinnitus severity with the Tinnitus Reaction Questionnaire.

Results

Type D tinnitus patients (n=94) did not differ significantly from non-Type D patients (n=171) on sociodemographic characteristics. Type D patients were significantly more anxious, depressed and vitally exhausted, and experienced more impaired HRQoL and increased tinnitus severity compared to non-Type D patients. Structural equation modelling showed that anxiety, depression and vital exhaustion had a direct influence on HRQoL and tinnitus severity, with a higher impact on mental HRQoL ($R^2=0.77$) compared to physical HRQoL ($R^2=0.33$). Vital exhaustion was a predictor of HRQoL and tinnitus severity, however, affected by enhanced levels of anxiety and depression. Type D personality directly increased symptoms of depression and anxiety, but not vital exhaustion. Type D was also a direct predictor of poor mental and physical HRQoL and high levels of tinnitus-related distress, although this influence worked mainly via depression and anxiety

Conclusions

All indicators of psychological distress were major predictors of poor HRQoL and tinnitus severity, in which the role of anxiety and depression was more enhanced compared to the role of vital exhaustion. The impact of Type D personality on HRQoL and tinnitus severity was largely mediated by anxiety and depression, although Type D also had a direct effect on these outcomes. Our results underline that for decreasing the implications of having tinnitus on HRQoL and disease-severity, treatment should be mainly directed at reducing anxiety and depression, especially in those patients with a Type D personality.

Introduction

Chronic subjective tinnitus is a common and often disturbing symptom, affecting 10-30% (1-3) of the adult population. About 1-5% of the population is severely affected by their tinnitus perception (2-4). Annoyance ratings and subjective loudness judgments frequently vary over time (5), and a wide variation exists in the impact of tinnitus on every day activities and health-related quality of life (HRQoL). It seems to bother some patients only slightly, whereas the impact on other patients is much more profound, leading to a reduction in daily activities and poor HRQoL (6-11).

Significantly perceived handicap and psychological distress have been described in about 40-50% of tinnitus sufferers (6). This is related to a variety of tinnitus-related complaints: anxiety, depression, insomnia, concentration problems, irritability, anger, frustration, inability to relax, feelings of helplessness, avoidance of noisy or quiet situations, withdrawal from social events, and emotional problems in relationships with family, friends and colleagues (7;11-21). The perceived distress due to tinnitus varies during the day and between days in most patients (22). Frequently a relation exists between the perception of severe tinnitus and the presence of excessive stress or emotions (23-27). Tinnitus noises become troublesome if the patient focuses the attention on the tinnitus (28), with tinnitus-related distress likely being modulated by its uncontrollability and maladaptive coping strategies (24;29-32). The negative impact of anxiety and depression on HRQoL has been described by several authors, using one measure or a combination of generic HRQoL and tinnitus-specific measures for disease severity (6-11;15;18;19;32-43).

Anxiety and depression are generally used as measures of psychological distress, symptoms generally being higher in tinnitus patients compared to controls (6-11;15;18;19;32-43). In this study we introduce vital exhaustion as an additional dimension of distress. Moreover, since perceived tinnitus severity was previously described to be associated with symptoms such as insomnia (14;44-46), concentration problems (7;46;47), and relaxation problems (48;49), we hypothesize tinnitus patients to suffer from increased fatigue and irritability. Vital exhaustion is a mental state that is defined as extreme fatigue, increased irritability and demoralization (50). Vital exhaustion has been measured predominantly in patients with cardiovascular disease (CVD) or cancer (51-56). In CVD patients, symptoms of fatigue and vital exhaustion have been described to be more prevalent in patients with a Type D (distressed) personality (57). Type D personality is characterized by negative affectivity and social inhibition. Therefore, Type D patients tend to experience increased negative emotions, generally feel sad and have a gloomy view of life, while not sharing these emotions with others due to fears of how they may react (58). The Type D construct was developed in cardiac patients and has been associated with poor HRQoL, increased levels of distress, less social support, and increased risk of mortality and morbidity in patients with CVD (59-61).

We previously described that tinnitus patients are more likely to have a Type D personality than healthy controls (62). Therefore, it seems likely that tinnitus patients with a Type D personality also experience more distress and poorer HRQoL.

In this study, we evaluated 1) the predictive power of psychological distress (i.e., vital exhaustion, anxiety and depression) on HRQoL and tinnitus severity in tinnitus patients with and without a Type D personality structure, and 2) whether psychological distress mediated the impact of Type D personality on HRQoL and disease severity in chronic tinnitus patients.

Materials and methods

Patients and design

Consecutive chronic subjective tinnitus sufferers (n = 265) seen at the Department of Otorhinolaryngology of the University Medical Center Groningen, the Netherlands, were included in the current study. Tinnitus patients aged ≥ 20 years were included, provided that they were consulting our clinic because of their tinnitus. They were all suffering from chronic tinnitus, defined as a duration of longer than three months. Tinnitus patients were excluded if tinnitus was not the main reason for consulting our clinic, or if they had objective tinnitus (determined by means of a diagnostic protocol for tinnitus) or chronic disease co-morbidity.

Procedure

Psychological distress, HRQoL, tinnitus severity, and Type D personality were measured by means of self-rating questionnaires filled in by the subjects in private. We chose self-rating questionnaires to avoid information bias (e.g. socially desirable responses) that might occur during personal interviews conducted by the otorhinolaryngologist in charge of the subject's medical examination and treatment. We asked subjects to answer the questionnaire at home and, in order to comply with the informed consent condition of protecting the patient's privacy, we also asked subjects to return their sealed questionnaires to the trial centre.

The study protocol was approved by the local medical ethics committee and all patients provided written informed consent.

Measures

Sociodemographic variables

Sociodemographic variables included age, gender, marital status, working status, and educational level.

Anxiety and Depression

The Hospital Anxiety and Depression Scale (HADS)

The HADS was used to assess anxiety and depressive symptoms (63;64). HADS is a 14-item measure, with 7 items contributing to the anxiety and depression subscales,

respectively. All items are answered on a 4-point Likert scale from 0 to 3 (score range of 0-21). The HADS is a reliable questionnaire that performs well in screening for the separate dimensions of anxiety and depression (65).

Vital exhaustion

Maastricht Questionnaire

Vital exhaustion, which is defined as mental fatigue, demoralisation and increased irritability, was assessed by the 21-item Maastricht Questionnaire (MQ) (66). Each item is rated according to a three-point scale (No = 0; ? =1; Yes =2), with a scale score obtained by summing the answers. Thus, the minimum score is 0 and the maximum 42, with a high score indicating a severe level of vital exhaustion. The MQ is an internally consistent measure, with Cronbach's alpha of 0.89.

Generic health-related quality of life (HRQoL)

The Short Form Health Survey 36 (SF-36)

The SF-36 is a valid, reliable, and widely used generic measure of HRQoL that has been used in a myriad of studies of patients with somatic disease and healthy populations, with the 36 items contributing to eight scales: physical functioning, role physical functioning, bodily pain, vitality, social functioning, role emotional functioning, mental health, and general health (67;68). These scales are standardized to a score from 0 (poor) to 100 (high) and can be combined into two component summary scores (i.e., a physical component summary (PCS) and mental component summary (MCS)). The SF-36 has good reliability estimates, with Cronbach's alpha ranging from .65 to .96 for all subscales (69).

Tinnitus Severity

Tinnitus Reaction Questionnaire (TRQ)

The Tinnitus Severity Questionnaire (TRQ), developed by Wilson in 1991 (39), was designed as a measure for distress associated with tinnitus. The TRQ consists of 26 questions that are answered on a 5-point Likert scale from 0 (not at all) to 4 (almost all of the time), with a score range of 0-104. A higher score indicates a higher level of perceived distress caused by tinnitus, also labelled as tinnitus severity. The test-retest reliability ($r = .88$) and internal consistency (Cronbach's alpha = .96) of the scale are good. A score on the TRQ correlates moderately to highly with clinician ratings ($r = .67$) and self-report measures of anxiety and depression ($r = .58-.87$) (39).

Type D (distressed) personality

DS14

Type D personality was assessed with the 14-item Type-D Scale (DS14) (58). The DS14 consists of the subscales negative affectivity (NA: 7 items; e.g., "I often feel unhappy") and social inhibition (SI: 7 items; e.g., "I am a 'closed' person"). Type D personality characterizes those who tend to experience increased negative emotions and who do not express these emotions in social interactions. A cut-off score ≥ 10 on

both subscales denotes those with a Type D personality (70). The DS14 has adequate reliability, with Cronbach's $\alpha = 0.89 / 0.88$ and 3-months test-retest reliability of $r = 0.72 / 0.82$ for the NA and the SI subscales, respectively (58).

Statistical analyses

Bivariate analyses

Discrete variables were compared using the chi-square test (Fisher's exact test when appropriate and the difference of proportions test (71)), and are presented as numbers and percentages. Continuous variables were normally distributed (Shapiro Wilk, $p > 0.05$), and were therefore compared with the Student t-test for independent samples and are presented as means \pm SD. Effect sizes (ES) were calculated only for the statistically significant results, since differences between groups that are due to random variation have no clinical relevance. Cohen's ES "d" for unrelated samples was used to estimate the magnitude of the difference between two groups (mean difference score/the pooled standard deviation). According to Cohen's thresholds, an ES of < 0.20 indicates a trivial difference, an ES of ≥ 0.20 to < 0.50 a small difference, an ES of ≥ 0.50 to < 0.80 a moderate difference and ≥ 0.80 a large difference (72). Middel et al. showed that $ES \geq 0.20$ reflects a clinically relevant difference (73) which was confirmed by Samsa et al. (74). Therefore, in this study an $ES \geq 0.20$ was considered to be a clinically relevant difference between Type D and non-Type D tinnitus patients. All statistical tests were two-tailed. A value of $p < 0.05$ was used for all tests to indicate statistical significance. All statistical analyses were performed using SPSS 13.0.1 for Windows.

Structural Equation Model (SEM)

A hypothesized model was investigated with structural equation modeling (SEM). SEM is a multivariable technique for testing the tenability of this model and is an elaboration of regression analysis. Path analysis was used to test a hypothesized model in which: (i) increased psychological distress (i.e., enhanced levels of depression, anxiety, and vital exhaustion) are predictors of poor mental and physical HRQoL and tinnitus severity, and (ii) Type D personality influences directly and indirectly mental and physical HRQoL and tinnitus severity through its effect on the indicators of psychological distress (i.e., anxiety, depression, and vital exhaustion). The latent construct Type D personality was estimated with the indicators of negative affectivity (NA) and social inhibition (SI), and defined as simultaneously high scores on both subscales (58). To allow for mutual comparisons between the path coefficients, the completely standardized solution was used. For judging the model fit, we used multiple criteria as suggested by Bentler and Bonett (75). The fit of the model was evaluated by means of (a) non-significant χ^2 indicating that a non-significant amount of variance in the data remains unexplained; a ratio of χ^2 to the degrees of freedom less than three generally indicates a good model fit (76); (b) the root mean square error of approximation (RMSEA) (77); (c) the standardized root mean square residual (SRMR) (78-80); (d) the comparative fit index (CFI)(81); and (e) the Adjusted Goodness of Fit Index (AGFI)(80). Both CFI and RMSEA were

used, as it has been argued that they provide more stable and accurate estimates than several of the other fit indices (81;82). Structural paths with related t-values > 1.96 can be regarded as significant at $p < 0.05$ (i.e., parameter estimates ± 1.96 standard errors should exclude 0). Given their complementary features, we used all five indices compared against their critical values to evaluate the models.

Results

Baseline characteristics stratified by Type D personality

Type D tinnitus patients ($n=94$) did not differ significantly from non-Type D tinnitus patients ($n=171$) on sociodemographic characteristics (Table 1).

Table 1. Sociodemographic characteristics of tinnitus patients stratified by Type D personality*

	Type D n= 94	Non-Type D n= 171	P-value	
Age (years)				
< 50 years	62 (66.0)	99 (57.9)	0.24 ²	
mean (SD)	55.68 (10.67)	55.77 (11.70)	0.95 ¹	
Males	69 (73.4)	116 (67.8)	0.41 ²	
Married/living with partner	81 (86.2)	153 (89.5)	0.43 ²	
Working	41 (43.6)	96 (56.8)	0.06 ²	
Educational level			95% CI	
elementary schooling	13 (15.1)	30 (18.6)	- 6.1	13.2 ³
lower schooling	20 (23.3)	31 (19.3)	- 14.8	6.8 ³
secondary schooling	21 (24.4)	43 (26.7)	- 9.1	13.7 ³
higher professional training	25 (29.1)	43 (26.7)	-14.1	9.4 ³
college education/university	7 (8.1)	14 (8.7)	- 6.7	7.8 ³

* Results are presented as n (%) unless otherwise indicated

¹. Student's T-test

². Fisher exact test

³. Difference of proportions test

Differences in distress and health-related quality of life between Type D and non-Type D tinnitus patients

Means, standard deviations, p-values and effect sizes for the different measures of psychological distress and HRQoL are shown in Table 2 . Type D tinnitus patients reported more anxiety, more depressive symptoms, more symptoms of vital exhaustion, and poorer HRQoL compared to non-Type D tinnitus patients. All differences in psychological distress and HRQoL outcomes were statistically significant ($p < 0.001$) and clinically relevant, as indicated by a large ES ($\geq .80$)

according to Cohen's effect size index. Nevertheless, the effect size of the difference in physical HRQoL between Type D and non-Type D patients was moderate (ES = 0.61) (Table 2).

Table 2. Differences in psychological distress and health-related quality of life stratified by Type D personality

	Type D (n = 94)		Non- Type D (n = 171)		p- value ¹	Effect Size (ES)	95% CI for ES	
	mean	SD	mean	SD			lower	Upper
psychological distress								
Anxiety	10.99	3.97	6.37	4.27	.0001	1.11	0.83	1.37
Depression	10.25	3.99	5.70	4.57	.0001	1.04	0.77	1.30
vital exhaustion	33.77	7.88	20.76	11.88	.0001	1.22	0.95	1.49
health-related QoL								
mental HRQoL	41.49	19.44	66.20	22.58	.0001	1.15	0.88	1.42
physical HRQoL	56.43	18.49	66.99	16.54	.0001	0.61	0.35	0.87
tinnitus severity	63.40	15.46	42.59	21.94	.0001	1.05	0.78	1.31

1. T-test

The relationship between Type D personality, psychological distress, health-related quality of life, and tinnitus severity (LISREL models)

All models were evaluated by examining the parameter estimates and by the indices of overall fit provided by LISREL. Residual correlations between NA and SI, between anxiety and depression were allowed as they belonged to the same measure and were assessed simultaneously. All model parameters met the criteria of good model fit as presented in Table 3. For each model, the Chi-square statistic indicated that a non-significant amount of variance in the data remained unexplained and relative to degrees of freedom was less than two, suggesting a good initial indicator of fit. The RMSEA was sufficiently low (<0.06), as was the SRMR (<0.005). The CFI value of 1.00 exceeded the .97 value and the AGFI index was > .95. Taken together, these results suggest that the hypothesized models fitted the data well.

Table 3. Parameters of model fit for mental, physical health-related quality of Life and tinnitus severity

Critical values of parameter estimates	χ^2	df	P	RMSEA	AGFI	SRMR	CFI [#]	R ²
				<0.06	>0.95	<0.05	>0.97	
parameter estimates								
Mental HRQoL (SF-36)	5.09	3	0.17	0.051	0.96	0.014	1.00	74
Physical HRQoL (SF-36)	5.82	3	0.12	0.06	0.95	0.019	1.00	33
Disease severity (TRQ)	5.96	3	0.11	0.56	0.95	0.001	1.00	65

the comparative fit index (CFI) with a value > .97 indicates a good fit. According to Hu and Bentler (1999), this criterion is more appropriate than the >.95 criterion, as the large number of severely misspecified models is unacceptable (81)

Figures 1-3 show the results of the path analysis, depicted by the direct and mediated paths between Type D personality, anxiety, depression and vital exhaustion, and the outcomes HRQoL and tinnitus severity, respectively. HRQoL is differentiated into components of mental and physical HRQoL.

Overall, as hypothesized in our model within our tinnitus population, the more anxious, depressed or vitally exhausted patients were, the poorer the perception of global mental HRQoL ($\beta = -.30$, $\beta = -.30$ and $\beta = -.29$, respectively) and physical HRQoL ($\beta = -.28$, $\beta = -.34$ and $\beta = -.45$, respectively), and the more increased tinnitus severity ($\beta = .21$, $\beta = .25$ and $\beta = .19$, respectively). These parameter estimates were significant (with negative betas for HRQoL and positive beta for TRQ), indicating that increased symptoms of anxiety, depression and vital exhaustion led to reduced mental and physical HRQoL and increased tinnitus severity. According to our hypothesis, Type D personality was associated with increased symptoms of anxiety, depression and vital exhaustion, and poorer HRQoL and increased tinnitus severity. Presented figures mainly demonstrated the mediating role of the three indicators of psychological distress on the impact of Type D on HRQoL and disease severity.




Figure 1. A path model of Type D personality as predictor of anxiety and depression as precursors of poor physical HRQoL in tinnitus patients. Coefficients are all statistically significant, * <0.05, ** <0.01

Figure 1: Physical HRQoL ($R^2 = 0.33$)

Anxiety, depression and vital exhaustion all turned out to predict physical HRQoL in the tinnitus population ($\beta = -.28, -.34, -.45$ respectively). Type D personality significantly influenced anxiety, depression and physical HRQoL ($\beta = .86, .84, .53$ respectively), but Type D did not directly influence vital exhaustion. However, the path analyses showed that the influence of Type D on vital exhaustion was present via anxiety ($\beta = 0.24$) and depression ($\beta = 0.33$). Overall, perceived physical HRQoL in the tinnitus population was predominantly determined by anxiety and depression, but also by Type D personality both by itself and via anxiety and depression.




Figure 2. A path model of Type D personality as predictor of anxiety and depression as precursors of poor mental HRQoL in tinnitus patients. Coefficients are all statistically significant, * <0.05 , ** <0.01

Figure 2: Mental HRQoL ($R^2=0.74$)

This model shows that all indicators of psychological distress predict mental HRQoL in the tinnitus population at a level comparable to that of physical HRQoL ($\beta=-0.30$, -0.30 , -0.29). Type D directly elevated anxiety ($\beta=0.87$) and depression ($\beta=0.85$) and decreased mental HRQoL ($\beta=-.26$), but did not increase vital exhaustion. The influence of Type D personality on vital exhaustion was only induced via anxiety ($\beta=0.25$) and depression ($\beta=0.32$).




Figure 3. A path model of Type D personality as predictor of anxiety and depression as precursors of increased disease severity in tinnitus patients. Coefficients are all statistically significant, * <0.05 , ** <0.01

Figure 3: Tinnitus severity: ($R^2 = 0.69$)

All direct paths from anxiety, depression and vital exhaustion exerted an effect on tinnitus severity. These parameter estimates were significant with positive betas indicating that more symptoms of anxiety, depression and vital exhaustion led to an increase in tinnitus severity ($\beta = 0.21$, $\beta = 0.25$ and $\beta = 0.19$, respectively). Type D directly influenced anxiety ($\beta = 0.86$), depression ($\beta = 0.84$), and tinnitus severity ($\beta = 0.25$), but no significant pathways were found between Type D and vital exhaustion. Type D mediated vital exhaustion via anxiety ($\beta = 0.24$) and depression ($\beta = 0.33$).

Predicting and mediating role of Type D across all models

Type D exerted a direct effect on anxiety, depression, mental and physical HRQoL, and tinnitus severity. Type D personality had a higher direct effect on physical HRQoL ($\beta = -0.53$, Figure 1) than on mental HRQoL ($\beta = -0.26$, Figure 2) and tinnitus severity ($\beta = 0.25$, Figure 3). Furthermore, Type D personality had a considerable direct effect on both anxiety and depression across all models, but did not have a direct effect on vital exhaustion in any of the models. According to all presented

figures, psychological distress (mainly anxiety and depression) mediated the relationship between Type D and HRQoL / tinnitus severity. Type D personality, anxiety and depression explained 55% of the variance in mental HRQoL and 31% of the variance in physical HRQoL and 65 % of the variance in tinnitus severity.

Discussion

In this study of a tinnitus population, we found statistically significant and clinically relevant differences in all measures of psychological distress (i.e., anxiety, depression, and vital exhaustion), HRQoL and disease severity between Type D versus non-Type D patients. The two personality types were comparable on sociodemographic characteristics. Since we were interested in the relative influence of the different indicators of psychological distress on mental and physical HRQoL and tinnitus severity, we performed path analyses by means of structural equation modeling (SEM) using LISREL . Anxiety and depression were direct major predictors of mental and physical HRQoL and tinnitus severity in the tinnitus population, but also indirect via enhancing vital exhaustion. Type D was a direct predictor of mental and physical HRQoL and tinnitus severity, but the impact of Type D on HRQoL and tinnitus severity was mainly mediated by anxiety and depression. Via strong influences on anxiety and depression, Type D only indirectly influenced vital exhaustion. These results provide evidence that increased levels of anxiety and depression are precursors of reduced mental and physical HRQoL and increased tinnitus severity, and that these indicators of psychological distress mediate the impact of Type D personality on HRQoL and disease severity in tinnitus patients.

In accordance to expectations, the presented LISREL-models showed that the explained variance in mental HRQoL was larger than for physical HRQoL. This indicates that psychological distress has a larger impact on mental HRQoL than on physical HRQoL in tinnitus patients. Although Type D was shown to exert a direct effect on mental and physical HRQoL and tinnitus severity, the majority of the effect of Type D on HRQoL and tinnitus severity was induced via the indicators of distress, and predominantly anxiety and depression. Since there is a large overlap between anxiety, depression, and mental HRQoL, this might explain the lower regression coefficient between Type D and mental HRQoL, with most of the variance following a pathway via anxiety and depression.

Psychological distress has been described to be the most important predictor of impaired HRQoL in chronic diseases in general (83-85). The term 'psychological distress' may accurately describe the individualized, subjective patient response to acute or chronic illness (83;85-93). If individuals experience psychological distress it may be manifested by an alteration from a stable baseline emotional state to one of anxiety, depression, demotivation, vital exhaustion, irritability, aggressiveness, self-depreciation, and even suicide in the extreme (50;84;90;94). Several tinnitus studies have reported that anxiety and depression are

important indicators of poor HRQoL contributing to the development of severe annoying tinnitus (11;14-18;24-27;31;36;95). Since vital exhaustion has previously been shown to be associated with psychological distress and poor HRQoL in patients with CVD (96-99), we decided to examine the role of vital exhaustion in tinnitus patients. Vital exhaustion was never previously investigated in tinnitus patients. Vital exhaustion was shown to directly predict both mental and physical HRQoL and tinnitus severity in tinnitus patients, and to exert an indirect effect on HRQoL via anxiety and depression. Our study demonstrated that the addition of vital exhaustion as an indicator for psychological distress may be worthwhile in the study of tinnitus patients.

The relation between Type D personality and different indicators of psychological distress is in accordance with previously performed studies in CVD patients (99;100). Our results demonstrated that Type D personality mainly influences anxiety and depression, and thereby indirectly vital exhaustion, HRQoL and tinnitus severity. This extends research on Type D personality, showing that the impact of Type D personality on HRQoL and disease severity is mainly mediated by psychological distress. In turn, this provides important information for both research and clinical practice, showing that tinnitus patients with a Type D personality are particularly prone to distress and therefore poor HRQoL and more tinnitus severity. Hence, in order to improve the HRQoL of tinnitus patients, interventions in these patients should be targeted towards reducing psychological distress, in particular in Type D patients.

In conclusion, we found that Type D tinnitus patients experienced more psychological distress and poorer HRQoL than non-Type D tinnitus patients. Anxiety, depression and vital exhaustion were direct predictors of all measures of HRQoL and disease severity in the tinnitus population. Although Type D was a direct predictor of mental and physical HRQoL and tinnitus severity, the impact of Type D on HRQoL and tinnitus severity was mainly mediated via anxiety and depression. These results provide evidence that increased levels of anxiety and depression are precursors of poor HRQoL and increased tinnitus severity, but that personality also plays an important role with its impact on HRQoL and disease severity being mediated by psychological distress. From a clinical point of view, these results underline that in order to improve HRQoL and decrease tinnitus severity in tinnitus patients, treatment and psychosocial interventions should be mainly directed at reducing anxiety and depression, especially in those patients with a Type D personality.

Reference List

- (1) Jastreboff PJ, Jastreboff MM. Tinnitus retraining therapy for patients with tinnitus and decreased sound tolerance. *Otolaryngol Clin North Am* 2003 Apr;36(2):321-36.
- (2) Axelsson A, Ringdahl A. Tinnitus--a study of its prevalence and characteristics. *Br J Audiol* 1989 Feb;23(1):53-62.
- (3) Heller AJ. Classification and epidemiology of tinnitus. *Otolaryngol Clin North Am* 2003 Apr;36(2):239-48.
- (4) Davis A, El Rafaie A. epidemiology of tinnitus. In: Singular Publishing Group, editor. *Tinnitus Handbook*. San Diego: 2000. p. 1-23.
- (5) Schechter MA, Henry JA. Assessment and treatment of tinnitus patients using a "masking approach." *J Am Acad Audiol* 2002 Nov;13(10):545-58.
- (6) Bauch CD, Lynn SG, Williams DE, Mellon MW, Weaver AL. Tinnitus impact: three different measurement tools. *J Am Acad Audiol* 2003 May;14(4):181-7.
- (7) Erlandsson SI, Hallberg LR. Prediction of quality of life in patients with tinnitus. *Br J Audiol* 2000 Feb;34(1):11-20.
- (8) Holgers KM, Zoger S, Svedlund K. Predictive factors for development of severe tinnitus suffering-further characterisation. *Int J Audiol* 2005 Oct;44(10):584-92.
- (9) Lynn SG, Bauch CD, Williams DE, Beatty CW, Mellon MW, Weaver AL. Psychologic profile of tinnitus patients using the scl-90-R and tinnitus handicap inventory. *Otol Neurotol* 2003 Nov;24(6):878-81.
- (10) Newman CW, Wharton JA, Jacobson GP. Self-focused and somatic attention in patients with tinnitus. *J Am Acad Audiol* 1997 Jun;8(3):143-9.
- (11) Scott B, Lindberg P. Psychological profile and somatic complaints between help-seeking and non-help-seeking tinnitus subjects. *Psychosomatics* 2000 Jul;41(4):347-52.
- (12) Baguley DM, Davies E, Hazell JW. A vision for tinnitus research. *Int J Audiol* 2003 Jan;42(1):2-3.
- (13) Tyler RS, Baker LJ. Difficulties experienced by tinnitus sufferers. *J Speech Hear Disord* 1983 May;48(2):150-4.
- (14) Dobie RA. Depression and tinnitus. *Otolaryngol Clin North Am* 2003 Apr;36(2):383-8.
- (15) Andersson G, Kaldo-Sandstrom V, Strom L, Stromgren T. Internet administration of the Hospital Anxiety and Depression Scale in a sample of tinnitus patients. *J Psychosom Res* 2003 Sep;55(3):259-62.
- (16) Andersson G, Stromgren T, Strom L, Lyttkens L. Randomized controlled trial of internet-based cognitive behavior therapy for distress associated with tinnitus. *Psychosom Med* 2002 Sep;64(5):810-6.
- (17) Andersson G. Psychological aspects of tinnitus and the application of cognitive-behavioral therapy. *Clin Psychol Rev* 2002 Sep;22(7):977-90.
- (18) Reynolds P, Gardner D, Lee R. Tinnitus and psychological morbidity: a cross-sectional study to investigate psychological morbidity in tinnitus patients and its relationship with severity of symptoms and illness perceptions. *Clin Otolaryngol Allied Sci* 2004 Dec;29(6):628-34.
- (19) Zoger S, Svedlund J, Holgers KM. Relationship between tinnitus severity and psychiatric disorders. *Psychosomatics* 2006 Jul;47(4):282-8.
- (20) Wilson PH. Classical conditioning as the basis for the effective treatment of tinnitus-related distress. *ORL J Otorhinolaryngol Relat Spec* 2006;68(1):6-11.
- (21) Folmer RL, Griest SE. Tinnitus and insomnia. *Am J Otolaryngol* 2000 Sep;21(5):287-93.
- (22) Andersson G, Lyttkens L, Larsen HC. Distinguishing levels of tinnitus distress. *Clin Otolaryngol Allied Sci* 1999 Sep;24(5):404-10.
- (23) Meric C, Gartner M, Collet L, Chery-Croze S. Psychopathological profile of tinnitus sufferers: evidence concerning the relationship between tinnitus features and impact on life. *Audiol Neurotol* 1998 Jul;3(4):240-52.
- (24) Budd RJ, Pugh R. Tinnitus coping style and its relationship to tinnitus severity and emotional distress. *J Psychosom Res* 1996 Oct;41(4):327-35.
- (25) Hallam RS, Jakes SC, Chambers C, Hinchcliffe R. A comparison of different methods for assessing the 'intensity' of tinnitus. *Acta Otolaryngol* 1985 May;99(5-6):501-8.
- (26) Collet L, Moussu MF, Dubreuil C, Disant F, Chanal JM, Morgon A. Psychological factors affecting outcome of treatment after transcutaneous electrotherapy for persistent tinnitus. *Arch Otorhinolaryngol* 1987;244(1):20-2.

- (27) Kirsch CA, Blanchard EB, Parnes SM. Psychological characteristics of individuals high and low in their ability to cope with tinnitus. *Psychosom Med* 1989 Mar;51(2):209-17.
- (28) Hallam RS, Jakes SC, Hinchcliffe R. Cognitive variables in tinnitus annoyance. *Br J Clin Psychol* 1988 Sep;27 (Pt 3):213-22.
- (29) Halford JB, Anderson SD. Tinnitus severity measured by a subjective scale, audiometry and clinical judgement. *J Laryngol Otol* 1991 Feb;105(2):89-93.
- (30) Vallianatou NG, Christodoulou P, Nestoros JN, Helidonis E. Audiologic and psychological profile of Greek patients with tinnitus--preliminary findings. *Am J Otolaryngol* 2001 Jan;22(1):33-7.
- (31) Zoger S, Svedlund J, Holgers KM. Psychiatric disorders in tinnitus patients without severe hearing impairment: 24 month follow-up of patients at an audiological clinic. *Audiology* 2001 May;40(3):133-40.
- (32) Zachariae R, Mirz F, Johansen LV, Andersen SE, Bjerring P, Pedersen CB. Reliability and validity of a Danish adaptation of the Tinnitus Handicap Inventory. *Scand Audiol* 2000;29(1):37-43.
- (33) Budd RJ, Pugh R. The relationship between locus of control, tinnitus severity, and emotional distress in a group of tinnitus sufferers. *J Psychosom Res* 1995 Nov;39(8):1015-8.
- (34) Andersson G, Vretblad P. Anxiety sensitivity in patients with chronic tinnitus. *Scand.J.of Behaviour Therapy* , 57-64. 2000.
- (35) Andersson G, Kaldo V, Stromgren T, Strom L. Are coping strategies really useful for the tinnitus patient? An investigation conducted via the internet. *Audiological medicine* 2, 54-59. 2004.
- (36) Holgers KM, Erlandsson SI, Barrenas ML. Predictive factors for the severity of tinnitus. *Audiology* 2000 Sep;39(5):284-91.
- (37) Meric C, Pham E, Chery-Croze S. Validation assessment of a French version of the tinnitus reaction questionnaire: a comparison between data from English and French versions. *J Speech Lang Hear Res* 2000 Feb;43(1):184-90.
- (38) Sullivan M, Katon W, Russo J, Dobie R, Sakai C. Coping and marital support as correlates of tinnitus disability. *Gen Hosp Psychiatry* 1994 Jul;16(4):259-66.
- (39) Wilson PH, Henry J, Bowen M, Haralambous G. Tinnitus reaction questionnaire: psychometric properties of a measure of distress associated with tinnitus. *J Speech Hear Res* 1991 Feb;34(1):197-201.
- (40) Andersson G, Carlbring P, Kaldo V, Strom L. Screening of psychiatric disorders via the Internet. A pilot study with tinnitus patients. *Nord J Psychiatry* 2004;58(4):287-91.
- (41) Harrop-Griffiths J, Katon W, Dobie R, Sakai C, Russo J. Chronic tinnitus: association with psychiatric diagnoses. *J Psychosom Res* 1987;31(5):613-21.
- (42) Langenbach M, Olderog M, Michel O, Albus C, Kohle K. Psychosocial and personality predictors of tinnitus-related distress. *Gen Hosp Psychiatry* 2005 Jan;27(1):73-7.
- (43) Rizzardo R, Savastano M, Maron MB, Mangialaio M, Salvadori L. Psychological distress in patients with tinnitus. *J Otolaryngol* 1998 Feb;27(1):21-5.
- (44) Asplund R. Sleepiness and sleep in elderly persons with tinnitus. *Arch Gerontol Geriatr* 2003 Sep;37(2):139-45.
- (45) Handscomb L. Use of bedside sound generators by patients with tinnitus-related sleeping difficulty: which sounds are preferred and why? *Acta Otolar.gol Suppl* 2006 Dec;(556):59-63.
- (46) Cronlein T, Langguth B, Geisler P, Hajak G. Tinnitus and insomnia. *Prog Brain Res* 2007;166:227-33.
- (47) Hallam RS, McKenna L, Shurlock L. Tinnitus impairs cognitive efficiency. *Int J Audiol* 2004 Apr;43(4):218-26.
- (48) Weber C, Arck P, Mazurek B, Klapp BF. Impact of a relaxation training on psychometric and immunologic parameters in tinnitus sufferers. *J Psychosom Res* 2002 Jan;52(1):29-33.
- (49) Andersson G. Psychological aspects of tinnitus and the application of cognitive-behavioral therapy. *Clin Psychol Rev* 2002 Sep;22(7):977-90.
- (50) Appels A, Mulder P. Fatigue and heart disease. The association between 'vital exhaustion' and past, present and future coronary heart disease. *J Psychosom Res* 1989;33(6):727-38.
- (51) Bergelt C, Christensen J, Prescott E, Gronbaek M, Koch U, Johansen C. Vital exhaustion and risk for cancer: a prospective cohort study on the association between depressive feelings, fatigue, and risk of cancer. *Cancer* 2005 Sep 15;104(6):1288-95.

-
- (52) Bages N, Appels A, Falger PR. Vital exhaustion as a risk factor of myocardial infarction: a case-control study in Venezuela. *Int J Behav Med* 1999;6(3):279-90.
 - (53) Casillas JM, Damak S, Chauvet-Gelinier JC, Deley G, Ornetti P. Fatigue in patients with cardiovascular disease. *Ann Readapt Med Phys* 2006 Jul;49(6):309-402.
 - (54) Appels A, van ET, Bar F, van der PG, Erdman RA, Assman M, et al. Effects of a behavioural intervention on quality of life and related variables in angioplasty patients: results of the EXhaustion Intervention Trial. *J Psychosom Res* 2006 Jul;61(1):1-7.
 - (55) Pedersen SS, Daemen J, van de SM, Sonnenschein K, Serruys PW, Erdman RA, et al. Type-D personality exerts a stable, adverse effect on vital exhaustion in PCI patients treated with paclitaxel-eluting stents. *J Psychosom Res* 2007 Apr;62(4):447-53.
 - (56) Pedersen SS, Denollet J, Daemen J, van de SM, de Jaegere PT, Serruys PW, et al. Fatigue, depressive symptoms, and hopelessness as predictors of adverse clinical events following percutaneous coronary intervention with paclitaxel-eluting stents. *J Psychosom Res* 2007 Apr;62(4):455-61.
 - (57) Pedersen SS, Denollet J. Type D personality, cardiac events, and impaired quality of life: a review. *Eur J Cardiovasc Prev Rehabil* 2003 Aug;10(4):241-8.
 - (58) Denollet J. DS14: Standard assessment of negative affectivity, social inhibition, and Type D personality. *Psychosomatic Medicine* 2005;67(1):89-97.
 - (59) Aquarius AE, Denollet J, Hamming JF, De Vries J. Role of disease status and type D personality in outcomes in patients with peripheral arterial disease. *American Journal of Cardiology* 2005;96(7):996-1001.
 - (60) Pedersen SS, Lemos PA, van Vooren PR, Liu TKK, Daemen J, Erdman RAM, et al. Type D personality predicts death or myocardial infarction after bare metal stent or sirolimus-eluting stent implantation - A Rapamycin-Eluting Stent Evaluated at Rotterdam Cardiology Hospital (RESEARCH) registry substudy. *Journal of the American College of Cardiology* 2004;44(5):997-1001.
 - (61) Pedersen SS, Denollet J, Ong A, van Domburg RT, Serruys PT. Lack of positive emotions and adverse prognosis post-PCI. *Psychology & Health* 2006;21:117-8.
 - (62) Bartels H, Middel JL, Pedersen SS, Staal MJ, Albers FW. The distressed (Type D) personality is independently associated with tinnitus adjusted for other personality characteristics: a case-control study. *Psychosomatics* Accepted for publication November 2007.
 - (63) Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983 Jun;67(6):361-70.
 - (64) Spinhoven PH, Ormel J, Sloekers PPA, Kempen GJIM. A validation study of the Hospital Anxiety and Depression Scale (HADS) in different groups of Dutch subjects. *Psychological Medicine* 1997;27(2):363-70.
 - (65) Bjelland I, Dahl AA, Haug TT, Neckelmann D. The validity of the Hospital Anxiety and Depression Scale. An updated literature review. *J Psychosom Res* 2002 Feb;52(2):69-77.
 - (66) Appels A, Hoppener P, Mulder P. A questionnaire to assess premonitory symptoms of myocardial infarction. *Int J Cardiol* 1987 Oct;17(1):15-24.
 - (67) Aaronson NK, Muller M, Cohen PD, Essink-Bot ML, Fekkes M, Sanderman R, et al. Translation, validation, and norming of the Dutch language version of the SF-36 Health Survey in community and chronic disease populations. *J Clin Epidemiol* 1998 Nov;51(11):1055-68.
 - (68) Wagner AK, Gandek B, Aaronson NK, Acquadro C, Alonso J, Apolone G, et al. Cross-Cultural Comparisons of the Content of SF-36 Translations across 10 Countries: Results from the IQOLA Project. *Journal of Clinical Epidemiology* 1998 Nov;51(11):925-32.
 - (69) Smith HJ, Taylor R, Mitchell A. A comparison of four quality of life instruments in cardiac patients: SF-36, QLI, QLMI, and SEIQoL. *Heart* 2000;84(4):390-4.
 - (70) Denollet J. Type D personality and vulnerability to chronic disease, impaired quality of life, and depressive symptoms. *Psychosom.Med.* 64:101. 2002.
 - (71) Newcombe RG, Altman DG. Proportions and their differences. In: Altman DG, Machin D, Bryant TN, Gardner MJ, editors. *Statistics with confidence*. Second ed. Bristol: British Medical Journal; 2005. p. 45-56.
 - (72) Cohen J. *Statistical power analysis for the behavioural sciences*. revised edition ed. New York: Academic Press; 1997.
 - (73) Middel B, Stewart R, Bouma J, van Sonderen E, van den Heuvel WJA. How to validate clinically important change in health-related functional status. Is the magnitude of the effect

- size consistently related to magnitude of change as indicated by a global question rating? *Journal of Evaluation in Clinical Practice* 2001;7(4):399-410.
- (74) Samsa G, Edelman D, Rothman ML, Williams GR, Lipscomb J, Matchar D. Determining clinically important differences in health status measures: a general approach with illustration to the Health Utilities Index Mark II. *Pharmacoeconomics* 1999 Feb;15(2):141-55.
- (75) Bentler PM, Bonett DG. Significance Tests and Goodness of Fit in the Analysis of Covariance-Structures. *Psychological Bulletin* 1980;88(3):588-606.
- (76) Kline RB. *Principles and Practice of Structural Equation Modeling*. New York: Guilford; 1998.
- (77) Browne MW, Cudeck R. Alternative ways of assessing model fit. In: Bollen KA, Long JS, editors. *Testing structural equation models*. Newbury Park (CA): Sage; 1993. p. 136-62.
- (78) Jaccard J, Wan CK. *LISREL approaches to interaction effects in multiple regression*. 1996. Thousand Oaks, CA, Sage. Quantitative Applications in the Social Sciences. Ref Type: Serial (Book, Monograph)
- (79) Jöreskog KG, Sörbom D. *LISREL 8.7 for Windows: User's reference guide*. Lincolnwood, IL: Scientific Software International, Inc; 2005.
- (80) Schermelleh-Engel K, Moosbrugger H, Müller H. Evaluating the fit of structural equation models: test of significance and descriptive goodness-of-fit measures. *Methods of Psychological Research Online* 2003;8(2):23-74.
- (81) Hu L, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis: conventional criteria versus new alternatives. *Structural Equation Modeling* 1999;6:1-55.
- (82) Maruyama GM. *Basics of structural equation modeling*. Thousand Oaks (CA): Sage; 1998.
- (83) Keles H, Ekici A, Ekici M, Bulcun E, Altinkaya V. Effect of chronic diseases and associated psychological distress on health-related quality of life. *Intern Med J* 2007 Jan;37(1):6-11.
- (84) Ridner SH. Psychological distress: concept analysis. *J Adv Nurs* 2004 Mar;45(5):536-45.
- (85) Noel PH, Williams JW, Jr., Unutzer J, Worchel J, Lee S, Cornell J, et al. Depression and comorbid illness in elderly primary care patients: impact on multiple domains of health status and well-being. *Ann Fam Med* 2004 Nov;2(6):555-62.
- (86) McClement SE, Woodgate RL, Degner L. Symptom distress in adult patients with cancer. *Cancer Nurs* 1997 Aug;20(4):236-43.
- (87) Holley S. Cancer-related fatigue. Suffering a different fatigue. *Cancer Pract* 2000 Mar;8(2):87-95.
- (88) Holley SK. Evaluating patient distress from cancer-related fatigue: an instrument development study. *Oncol Nurs Forum* 2000 Oct;27(9):1425-31.
- (89) May M, McCarron P, Stansfeld S, Ben-Shlomo Y, Gallacher J, Yarnell J, et al. Does psychological distress predict the risk of ischemic stroke and transient ischemic attack? The Caerphilly Study. *Stroke* 2002 Jan;33(1):7-12.
- (90) Nordin K, Liden A, Hansson M, Rosenquist R, Berglund G. Coping style, psychological distress, risk perception, and satisfaction in subjects attending genetic counselling for hereditary cancer. *J Med Genet* 2002 Sep;39(9):689-94.
- (91) Shaver JL, Johnston SK, Lentz MJ, Landis CA. Stress exposure, psychological distress, and physiological stress activation in midlife women with insomnia. *Psychosom Med* 2002 Sep;64(5):793-802.
- (92) Rustoen T, Wahl AK, Hanestad BR, Lerdal A, Paul S, Miaskowski C. Gender differences in chronic pain--findings from a population-based study of Norwegian adults. *Pain Manag Nurs* 2004 Sep;5(3):105-17.
- (93) Yates WR, Mitchell J, Rush AJ, Trivedi MH, Wisniewski SR, Warden D, et al. Clinical features of depressed outpatients with and without co-occurring general medical conditions in STAR*D. *Gen Hosp Psychiatry* 2004 Nov;26(6):421-9.
- (94) Massee R. Qualitative and quantitative analyses of psychological distress: methodological complementarity and ontological incommensurability. *Qualitative Health Research* 10, 411-423. 2000.
- (95) Simpson RB, Nedzelski JM, Barber HO, Thomas MR. Psychiatric diagnoses in patients with psychogenic dizziness or severe tinnitus. *J Otolaryngol* 1988 Oct;17(6):325-30.
- (96) Appels A, Falger PR, Schouten EG. Vital exhaustion as risk indicator for myocardial infarction in women. *J Psychosom Res* 1993 Dec;37(8):881-90.

-
- (97) Appels A, Kop W, Bar F, de SH, Mendes de LC. Vital exhaustion, extent of atherosclerosis, and the clinical course after successful percutaneous transluminal coronary angioplasty. *Eur Heart J* 1995 Dec;16(12):1880-5.
 - (98) Kop WJ, Appels AP, Mendes de Leon CF, de Swart HB, Bar FW. Vital exhaustion predicts new cardiac events after successful coronary angioplasty. *Psychosom Med* 1994 Jul;56(4):281-7.
 - (99) Pedersen SS, Denollet J, Daemen J, van de SM, de Jaegere PT, Serruys PW, et al. Fatigue, depressive symptoms, and hopelessness as predictors of adverse clinical events following percutaneous coronary intervention with paclitaxel-eluting stents. *J Psychosom Res* 2007 Apr;62(4):455-61.
 - (100) Pedersen SS, Middel B. Increased vital exhaustion among type-D patients with ischemic heart disease. *J Psychosom Res* 2001 Aug;51(2):443-9.
